


ExamsRoad.com



Inorganic Chemistry

H₂ Hydrogen

- Group → 1 (Alkali metals) Period → 1 Non metal
- Hydrogen shows similar characteristics to both alkali metals (Gr. 1) and halogens (Gr. 17)
- Dihydrogen (H₂): Most abundant molecule in the universe (70%)
- Isotopes: protium (¹H), deuterium (²H), tritium (³H)
- most abundant
- 0.0156% in terrestrial Hydrogen, mostly in form of HD
- only radioactive emits β particle t_{1/2} = 12.33 years

PREPARATION

- Granulated Zn and dilute HCl: $\text{Zn} + 2\text{HCl} \rightarrow \text{ZnCl}_2 + \text{H}_2$ [Lab prep] [Mina Aschido]
- Zn and aqueous alkali: $\text{Zn} + 2\text{NaOH} \rightarrow \text{Na}_2\text{ZnO}_2 + \text{H}_2$
- Electrolysis of acidified water: $2\text{H}_2\text{O}(\text{l}) \xrightarrow{\text{electricity}} 2\text{H}_2(\text{g}) + \text{O}_2(\text{g})$ [Underwater Kaminari]
- High purity (>99.95%) H₂ is obtained by electrolyzing warm aqueous barium hydroxide solution w/ nickel electrodes.
- Electrolysis of brine solution: $2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$ [As a byproduct]
- Steam on hydrocarbon/coke at high T & catalyst: $\text{C}_n\text{H}_{2n+2} + n\text{H}_2\text{O} \xrightarrow[\text{Ni}]{1270\text{K}} n\text{CO} + (2n+1)\text{H}_2$
- $\text{CH}_4(\text{g}) + \text{H}_2\text{O}(\text{g}) \xrightarrow[\text{Ni}]{1270\text{K}} \text{CO}(\text{g}) + 3\text{H}_2(\text{g})$ → water gas / synthesis gas / syngas

Prep of Syngas from Coal (Coal Gasification):




PROPERTIES

Physical






- Colourless, Odourless, tasteless
- Combustible, insoluble in H₂O
- Lighter than air,

Chemical

- (H-H bond dissociation enthalpy is the highest for single bond between two atoms of any element.)
- Can lose 1 e⁻ to form H⁺, gain to form H⁻ or form covalent bond.




- Reaction with F occurs even in dark
- Iodine needs catalyst.



Reaction with Organic compounds

- Hydrogenation of vegetable oils using nickel as catalyst gives edible fats.
- Hydroformylation of olefins yields aldehydes which further undergo reduction to give alcohols.



Uses of Hydrogen

- Synthesis of ammonia
- manufacture of vanaspathi fat
- manufacture of organic compounds particularly methanol
- manufacture of metal hydrides.
- preparation of HCl
- Reducing heavy metal oxides to metal
- Atomic H and oxy-hydrogen torches used for cutting and welding.
- [Atomic H are allowed to recombine on the surface generating 4000K]
- Rocket fuel.
- Used in fuel cells for generating electrical energy


Uses of H₂

Amelia takes a fat organo-metallic pipe to weld the rocket fuel tank lifted by hydrolie, but the electricity read out



Hydrides (E H_x → E⁺ + xH⁻)

→ Ionic or Saline Hydrides • stoichiometric compounds with highly electropositive elements

- Generally ionic but covalent character found in light metal hydrides.
- BaH₂, MgH₂ are polymeric
- Ionic ones are crystalline, non-volatile, non-conducting in solid state.
- But melts of ionic ones conduct electricity and liberate H₂ gas.
- Saline hydrides react violently with H₂O producing H₂ gas.



- LiH is rather unreactive at moderate temperatures with O₂ or Cl₂.
- It is therefore, used in the synthesis of other useful hydrides.



Prep of Hydrogen

Mina walked in the lab with a piece of Zinc to make H₂ and found Kaminari swimming in acidified water. After drying him for high purity bakugo and tetsu tetsu were called. They took him to Doctor Brown. There was steam coming out of his room. In such hotness he was drinking coke with his cat in lap. [MP]


● Covalent / Molecular Hydrides • covalent molecules formed with the p block elements. [CH_4 , NH_3 , H_2O , HF etc.]
 • Being covalent, they are volatile

- electron-deficient: Incomplete octate, electron acceptors, acts as Lewis acid, group (13)
- electron-precise: Complete octate, All group 14 hydrides
- electron-rich: More than 8 e⁻ in last shell, group (15-17), electron donor, Lewis base

● Metallic / Non-stoichiometric Hydrides • compounds with d block and f block [7, 8, 9th Group elements don't]
 • from 6th group only Chromium forms CrH • Good conductors of heat and electricity. $\text{PdH}_{0.6-0.9}$
 • Almost always non-stoichiometric e.g. $\text{LaH}_{2.87}$, $\text{YbH}_{2.55}$, $\text{TiH}_{1.5-1.8}$, $\text{ZrH}_{1.3-1.75}$, $\text{VH}_{0.56}$, $\text{NiH}_{0.6-0.7}$
 • Some metals absorb H_2 into their lattice, they are called interstitial hydrides. This property is used is widely used in catalytic reduction / hydrogenation.
 • Pd, Pt can accommodate a very large volume of Hydrogen

● WATER • Human body has about 65%.


PROPS




- highly polar molecule • sp^3 hybridised
- In liquid phase, there exist intermolecular H-bond.
- Density of ice is less than water.

Chemical



• Amphoteric Nature: Reacts as an acid with NH_3 and as a base with H_2S .



• Auto-protolysis (self-ionization)





• Hydrolysis Reaction Due to high dielectric constant, it has a very strong hydrating tendency.



● Hard Water Contains ~~calcium~~ bicarbonates, and chlorides and sulphates. Doesn't readily form lather. (forms scum/precipitate with soap)

[Soap containing sodium stearate ($\text{C}_{17}\text{H}_{35}\text{COONa}$) reacts with hard water to precipitate out Ca/Mg stearate]





Classification → ① Temporary Hardness
 ② Permanent Hardness

● Permanent Hardness [Due to Mg/Ca chlorides/sulphates]

can be removed by

• treating w/ washing soda (Na_2CO_3) Converts the soluble salts into insoluble carbonates.



Physical

- tasteless, colourless liquid
- high mp freezing point, b.p, heat of vap, heat of fusion
- high thermal conductivity, surface tension, high specific heat capacity
- dipole moment and dielectric const.
- Anomalous behaviour


Structure of Ice

Each O atom is surrounded tetrahedrally by four O atoms.

• Redox involving H_2O H_2O is easily reduced to H_2 by highly electropositive metals.



Oxidised to O_2 during photosynthesis and with F_2 too.



• Hydrates Formation From aqueous solution many salts can be crystallised as hydrated salts. Classification →

→ coordinated water $[\text{Cr}(\text{H}_2\text{O})_6]^{3+} 3\text{Cl}^-$

→ interstitial water $\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$

→ Hydrogen-bonded water $[\text{Al}(\text{H}_2\text{O})_4]^{2+} \text{SO}_4^{2-}$, $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$



● Soft Water: Basically pure from minerals and readily forms lather with soap.

→ Temporary Hardness [Due to Mg/Ca bicarbonate]



can be removed by

• Boiling By boiling, soluble $\text{Mg}(\text{HCO}_3)_2$ and $\text{Ca}(\text{HCO}_3)_2$ are converted into insoluble $\text{Mg}(\text{OH})_2$ and $\text{Ca}(\text{HCO}_3)_2$ CaCO_3 respectively, which forms precipitate.

• Clark's Method Calculated amount of added lime precipitates out CaCO_3 and $\text{Mg}(\text{OH})_2$




• Calgon's method Sodium hexameta phosphate ($\text{Na}_6\text{P}_6\text{O}_{18}$) commercially called 'calgon'. The following reactions occur.




↑ This complex anion keeps the Mg^{2+} and Ca^{2+} ions in solution

● Ion-exchange method: This method is also known as Zeolite / Permutit process. Hydrated sodium aluminium silicate is zeolite / permutit $[(NaAlSi_3O_8)_n \cdot nH_2O]$ can be written as NaZ




When all the Na is used, it is regenerated by NaCl






● Synthetic Resins Method: Ion exchange resin (RSO_3H) is changed to RNa by NaCl. It exchanges Na^+ with Ca^{2+} and Mg^{2+}



Pure de-mineralised (de-ionized) water free from all soluble mineral salts is obtained by passing water successively through a cation exchange (H^+) and an anion exchange (OH^-) resins.




In this process, water is acidified, in anion exchange, it gets neutralised.




● Hydrogen Peroxide (H_2O_2) $[H-O-O-H]$


● PREP ● Acidifying Barium Peroxide and removing water by evaporation under low press.




Used for making D_2O_2 ● Peroxodi sulphate, obtained by electrolytic oxidation of acidified sulphate at high current density on hydrolysis




● INDUSTRIALLY, prepared by Auto-oxidation of 2-alkylanthraquinols.



● Structure



● Gas Phase







● Solid phase

● Physical Properties

- colourless (very pale blue)
- forms a hydrate ($H_2O_2 \cdot H_2O$)
- 30% solⁿ is marketed as 100 volume.

● Chemical properties [Acts as both oxidising & reducing agent]



● Storage [Decomposes slowly on the exposure of light] $2H_2O_2(l) \longrightarrow 2H_2O(l) + O_2(g)$

→ In presence of metal or traces of alkali, the above reaction is catalysed. It is stored in wax-lined glass or plastic vessels in dark. Urea can be added as stabiliser. Kept away from dust as it causes explosive decomposition.

● Uses → Used as hair bleach and mild disinfectant. Sold as an antiseptic with the name perhydro/


→ manufacture chemicals like sodium perborate and per carbonates (detergents)

→ synthesis of hydroquinone, tartaric acid, cephalosporin etc

→ bleaching agent for textile, paper pulp, leather, oils, fats etc

→ in pollution control, industrial effluents, oxidation of cyanides, restoration of aerobic conditions to sewage

● Heavy Water (D_2O) → used as moderator in nuclear reactors, used in preparation of other deuterium.



● H_2 as a fuel

S-Block

● Groups 1 & 2 ● Group 1: Li, Na, K, Rb, Cs, Fr $[ns^1]$ ● Group 2: Be, Mg, Ca, Sr, Ba, Ra $[ns^2]$
 ↖ Alkali metals ↖ Alkaline earth metals

- Francium is radioactive, ^{223}Fr has a $t_{1/2} = 21$ minutes
- Beryllium (Be) is rare (only $10^{-10}\%$ of igneous rocks)
- Li shows similarities to Mg and Be to Al [Diagonal relationship] \rightarrow Due to similar size and ρ_m ratio.

Group 1: Alkali metals • Highly electropositive metals and lose 1 electron to form M^+ ions.

|| TRENDS ||

II TRENDS II
 → Atomic/Ionic Radii: Increase down the group $[Li < Na < K < Rb < Cs < Fr]$. Ions are smaller than parent atom.
 down the group $[Li > Na > K > Rb > Cs > Fr]$ ● Lithium has max hydration enthalpy, thus Li salts are mostly hydrated $[LiCl \cdot 2H_2O]$

→ Atomic / Ionic Radii: Increase down the group $[Li > Na > K > Rb > Cs > Fr]$

→ Hydration Enthalpy: Decrease down the group $[Li > Na > K > Rb > Cs > Fr]$

PHYSICAL PROPERTIES

● All are silvery white, soft and light metals. ● Density ~~decreases~~^{increases} from Li to Cs [Exception, K is lighter than Na]

② M.P and B.P.s are low due to weak metallic bonding.

- ⊗ M.P and B.P.s are low due to weak metallic bonding.
- ⊗ Impart characteristic flame in oxidising flame due to shifting of excited electron and coming back.


<u>Metal</u>	<u>Colour</u>	<u>λ (nm)</u>
Li	Crimson Red	670.8
Na	Yellow	589.2
K	Violet	766.5
Rb	Red violet	780.0
Cs	Blue	455.5

II CHEMICAL PROPERTIES II

II CHEMICAL PROPERTIES II

→ Reaction with Air: They tarnish in dry air due to formation of oxides, which in turn reacts with moisture to form hydroxide. Na forms peroxide, Li forms monoxide, others form superoxides [O_2^- is stable w/ large cations (K, Rb, Cs)]

$2M + O_2 \rightarrow 2MO_2$ (superoxide) [$M = K, Rb, Cs$]



$4\text{Li} + \text{O}_2 \longrightarrow 2\text{Li}_2\text{O}$ (oxide) / $2\text{Na} + \text{O}_2 \longrightarrow \text{Na}_2\text{O}_2$ (peroxide)
 • Li shows exceptional behaviour by reacting directly with N_2 of air to form nitride $[\text{Li}_3\text{N}]$
 • These metals are kept submerged in kerosene oil.

- Due to high reactivity towards air, these metals are kept submerged in kerosene oil.
- Some react with water to form hydroxide and dihydrogen. $[2M + 2H_2O \rightarrow 2M^+ + 2OH^- + H_2]$

→ Reaction with Water: Alkali metals react with water to form hydroxide and hydrogen gas. Reaction of Li with H_2O is less vigorous than Na.

Li \rightarrow most $-ve$ E^\ominus value, Na \rightarrow least $-ve$ E^\ominus value, Reaction of Li with H_2O is explosive, This is due to small size and very high Hydration energy, Other metals react explosively, Li reacts with NH_3 to form solvated species $Li(NH_3)_4^+$ and $Li(NH_3)_2^-$.

Also react with proton donors like alcohol, gaseous ammonia and terminal alkynes.

Also react with proton donors like alcohol, gaseous ammonia and H_2 .
 → Reaction with H_2 : They react with H_2 at about 673 K (Li at 1073 K) to form hydrides. [All ionising w/ high melting point]
 Due to high Polarising power of Li^+ (Fajan's Rule)



Reaction with Halogens: $2M + H_2 \longrightarrow 2M^+H^-$


→ Reaction with Halogens: React vigorously to form ionic halides. @ Exception: Li-halides are somewhat covalent [LiI is most covalent due to large anion]

$M(s) \longrightarrow M(g)$ [Sublimation enthalpy]

→ Reaction with Halogens: ⦿ React vigorously to form ionic
 $M(s) \rightarrow M(g)$ [Sublimation enthalpy]
 ⦿ Reaction with H_2 to form ionic hydrides. $M(g) \rightarrow M^+(g) + e^-$ [Ionisation enthalpy]

→ Reducing Nature: Strong reducing agents, Li strongest, Na weakest. $M(s) \rightarrow M(g)$ [Atomisation enthalpy]
 $M(g) \rightarrow M^+(g) + e^-$ [Ionisation enthalpy]
 $M^+(g) + H_2O \rightarrow M^+(aq)$ [Hydration enthalpy]

→ Solutions in liquid ammonia:
 1. Dissolve in liq. NH_3 giving deep blue solution which are conducting in nature.
 2. The blue colour is due to the ammoniated e^- which absorbs energy and
 3. Solution is paramagnetic.



$M + (x+y)NH_3 \longrightarrow [M(NH_3)_x]^+ + [e(NH_3)_y]^-$ The blue colour is due to the ammoniated e^- which imparts blue colour to the solution. Solution is paramagnetic.

On standing slowly, liberates hydrogen resulting in formation of amide. $[M^+_{(am)} + e^- + NH_3(l) \longrightarrow MNH_2(am) + \frac{1}{2}H_2(g)]$

In concentrated solution, the blue colour changes to bronze colour and becomes diamagnetic.

- On standing slowly, liberates hydrogen resulting in formation of amide. $[M]^{+}_{(aq)} + e^{-} + NH_3(l) \rightarrow [M]^{+}_{(aq)} + NH_3(l)$
- In concentrated solution, the blue colour changes to bronze colour and becomes diamagnetic.

USES

USES

- Li with Pb makes 'white metal' ball bearings for motor engine.
- Li with Al is used to make aircraft parts.
- Also ^(Li) used to make electrochemical cells.

- Li with Pb makes 'white metal' ball bearings for missiles.
- Li with Mg alloy is used to make armour plates.
- Also (Li) used to make electrochemical cells.
- Pb, Fe, and Pb-Mg \rightarrow used to be used

- Li with Pb makes white metal ball bearings.
- Li with Mg alloy is used to make armour plates. Also ^(Li) used to make electrochemical cells.
- Na is used to make Na/Pb alloy needed to make $PbEt_4$ and $PbMe_4$. \rightarrow used to be used as anti-knock additive.
- Na is used to make Na/Pb alloy needed to make $PbEt_4$ and $PbMe_4$.
- K has vital role for biological systems. $KCl \rightarrow$ fertilizer.

- Na is used to make Na/Pb alloy needed to make $PbEt_4$ and $PbCl_4$.
- liquid Na is used as coolant in fast breeder nuclear reactors.
- K has vital role for biological systems.
- KCl \rightarrow fertilizer


KOH is used in the manufacture of soft soaps. Also used as an excellent absorbent of CO_2 .

• Cs is used in devising photoelectric cells.

COMPOUNDS OF ALKALI METAL

→ Oxides & Hydroxides • Li forms oxide (Li_2O) and peroxide (Li_2O_2), Na forms peroxide (Na_2O_2) and superoxide (NaO_2), others form only superoxides.

• Increasing stability of peroxide and hydroxide, as the size of the metal ion increases, is due to the stabilisation of large anions by larger cations through lattice energy effects.
 formation of hydroxide by hydrolysis of oxide



- The oxides and peroxides are colourless when pure, but the superoxides are yellow or orange in colour.
- Superoxides are paramagnetic (O_2^-). Hydroxides made for oxides and H_2O are all white crystalline solids.
- Alkali metal hydroxides are strongest of all bases and dissolve freely in H_2O , releases heat on intense hydration.

→ Halides • Crystalline colourless solids with high melting point.

• All these halides have high negative enthalpies of formation; $\Delta_f H^\ominus$ values for fluorides become less negative as we go down the group but that for chlorides, bromides and iodides are reverse.

• For a given metal $\Delta_f H^\ominus$ always becomes less negative from fluoride to iodide

• m.p & b.p → Fluoride > Chloride > Bromide > Iodide → All are soluble in water.

• **EXC** Low solubility of LiF in water is due to its high lattice enthalpy whereas low solubility of CsI is due to smaller hydration enthalpy of its two ions.

• Other halides of Li are soluble in ethanol, acetone and ethylacetate; LiCl is soluble in pyridine also.

→ Salts of Oxo-Acids • Acidic proton and oxo group both are attached to the same hydroxyl group.
 $\text{H}_2\text{CO}_3 \rightarrow \text{O}=\text{C}(\text{OH})_2$
 $\text{H}_2\text{SiO}_4 \rightarrow \text{O}_2\text{Si}(\text{OH})_2$

• Alkali metals form salts with all the oxo acids (Generally soluble).

• Carbonates and bicarbonates are highly stable to heat. • As electropositivity increase down the group, stability increases.

• Li_2CO_3 is not stable, being small cation it polarises CO_3^{2-} ion leading to formation of more stable Li_2O and CO_2

• LiHCO_3 doesn't exist as a solid.

Anomalous Properties of Li

→ Cause: • exceptionally small size. • High polarising power (Eq) (e/r ratio).

→ Points: • Increased covalent character which results in solubility in organic compounds. • Diagonal relationship with Mg.

• harder and higher m.p., b.p. than other alkali metals • Li is least reactive • strongest reducing agent among Group 1

• Forms nitride on combustion like no other metal, alkali metal.

• LiCl is deliquescent and crystallises as a hydrate ($\text{LiCl} \cdot 2\text{H}_2\text{O}$), other alkali metal chlorides don't

• LiHCO_3 is not obtained in the solid form while all other elements form solid hydrogen carbonate.

• Li unlike other alkali metals don't form ethynide on reaction with ethyne.

• LiNO_3 gives Li_2O on heating, other decompose $[4\text{LiNO}_3 \rightarrow 2\text{Li}_2\text{O} + 4\text{NO}_2 + \text{O}_2 / 2\text{NaNO}_3 \rightarrow 2\text{NaNO}_2 + \text{O}_2]$

• LiF and Li_2O are much less soluble due to covalency.

Similarities between Li and Mg

• Both Li and Mg are harder, and lighter than other elements in respective group.

• Both react slowly with water. Oxides and hydroxides are much less soluble. hydroxides decompose on heating.

• The oxides as Li_2O and MgO do not combine with excess O_2


• Both Li_2O are soluble in ethanol

• Both LiCl and MgCl_2 are deliquescent and crystalline.

COMPOUNDS OF Na


→ Solvay Soda ($\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$) • Generally prepared by solvay process.

• Advantage of low solubility of NaHCO_3 is taken, as, it gets precipitated in the reaction of NaCl and $(\text{NH}_4)\text{HCO}_3$. $(\text{NH}_4)\text{HCO}_3$ is created by passing CO_2 to a conc solution of NaCl saturated with NH_3



• NaHCO_3 crystal separates. These are heated to give Na_2CO_3 $[2\text{NaHCO}_3 \rightarrow \text{Na}_2\text{CO}_3 + \text{CO}_2 + \text{H}_2\text{O}]$

• In this process NH_3 is recovered when the solution containing NH_4Cl is treated with $\text{Ca}(\text{OH})_2$.



• Solvay process can't be extended to the manufacture of K_2CO_3 bcz KHCO_3 is too soluble to be precipitated by the addition of NH_4HCO_3 to a saturated solution of KCl .

→ Sodium Chloride (NaCl) Most abundant source is sea water containing 2.7 to 2.9% by mass of NaCl.

- In tropical countries, it is prepared by solar evaporation [50 lakh tons/year in India]
- Crude sodium chloride, generally obtained from crystallisation of brine solution, contains Na_2SO_4 , CaSO_4 , CaCl_2 , MgCl_2 as impurities.
- CaCl_2 , MgCl_2 are impurities because they are deliquescent (absorbs moisture)
- To obtain pure NaCl, crude salt is dissolved in minimum amount of water and filtered to remove insoluble impurities. The solution is then saturated with HCl gas. Crystals of pure sodium chloride separate out. Calcium and MgCl_2 , being more soluble, remains in solution.
- Sodium chloride melts at 1081 K, solubility = 36g, solubility doesn't increase appreciably with increase in temp.

Uses: Used as common salt Used in preparation of Na_2O_2 , NaOH and Na_2CO_3

→ Caustic Soda (NaOH) prepared commercially by electrolysis of NaCl in Carter-Kellner cell.



Na and Hg forms amalgam at cathode
 Cl_2 is evolved at anode

- Treated with water, $2\text{Na-amalgam} + 2\text{H}_2\text{O} \longrightarrow 2\text{NaOH} + 2\text{H}_2 + \text{H}_2$
- NaOH is white translucent solid, m.p. = 591 K, readily soluble in water to give strong alkaline solution.
- Crystals of NaOH are deliquescent, NaOH at the surface reacts with CO_2 in atmosphere to form Na_2CO_3 .

Uses: Manufacture of Soap, paper, artificial silk in petroleum refining Purification of bauxite
In textile industries for mercerising cotton fabrics for preparation of purest acids As a laboratory reagent

→ Baking Soda (NaHCO_3) Decomposes on heating to release CO_2 making cakes or pastries fluffy.

- Made by saturating a solution of Na_2CO_3 with CO_2 , white crystalline NaHCO_3 powder, being less soluble, separates out.
 $\text{Na}_2\text{CO}_3 + \text{H}_2\text{O} + \text{CO}_2 \longrightarrow 2\text{NaHCO}_3$
- Mild antiseptic and used in fire extinguishers.

Biological Importance of Na and K

- Na ions are found primarily on the outside of cells, being located in blood plasma and in the interstitial fluid which surrounds the cells. These ions participate in the transmission of nerve signals, in regulating the flow of water across cell membranes and in the transport of sugars and amino acids into cells.
- K^+ ions are the most abundant cations within cell fluids, where they activate enzymes, participate in the oxidation of glucose to produce ATP and with Na^+ , are responsible for the transmissions of nerve signals.
- The ionic gradient across various blood cells demonstrate that a discriminatory mechanism called the sodium-potassium pump, operates across the cell membranes which consumes more than one-third of the ATP used by resting animal and about 15 kg per 24 h in human.

Group-2

Electronic conf: ns^2 Elements: Be, Mg, Ca, Sr, Ba, Ra

TRENDS → Atomic Radii/Ionic Radii: $\text{Be} < \text{Mg} < \text{Ca} < \text{Sr} < \text{Ba} < \text{Ra}$ [All are smaller than corresponding Alkali metals]

→ IE: Ionisation Enthalpy decreases down the group. [$\text{Ra} > \text{Ba}$]

→ Hydration Enthalpies: $\text{Be}^{2+} > \text{Mg}^{2+} > \text{Ca}^{2+} > \text{Sr}^{2+} > \text{Ba}^{2+}$ [More than alkali metals] Thus their salts are generally hydrated unlike alkali metal

PHYSICAL PROPERTIES: Generally are silvery white, lustrous and relatively soft but harder than alkali metals.

- Be, Mg are greyish M.Ps and B.Ps are higher than corresponding alkali metals due to smaller size.
- Electropositive character increases down the group from Be to Ba
- Ca, Sr, Ba impart characteristic brick red, crimson and apple green colours respectively to the flame. The e- in Be and Mg are too tightly held to be excited and then release energy, hence no colour.
- High conductivity and typical metal properties.

CHEMICAL PROPERTIES [Less reactive than alkali metals / Reactivity increases down the group]

→ Rxn w/ Air and H_2O : Be, Mg are kinetically inert to air and water due to formation of oxide film on their surface.

- But powdered Be burns brilliantly on ignition in air to give BeO and Be_3N_2
- Mg is more electropositive and burns with dazzling brilliance in air to give MgO and Mg_3N_2 .
- Ca, Sr, Ba are readily attacked by air to form oxide and nitride. React also with H_2O with increasing vigour even in cold to form hydroxides.

→ Rxn w/ Halogen: $[\text{M} + \text{X}_2 \longrightarrow \text{MX}_2]$ But Thermal decomposition of $(\text{NH}_4)_2\text{BeF}_4$ is the best route for the preparation of BeF_2 , and BeCl_2 is conveniently made from the oxide. $[\text{BeO} + \text{C} + \text{Cl}_2 \xrightarrow{600-800\text{K}} \text{BeCl}_2 + \text{CO}]$

→ Rxn w/ H_2 : All except Be combine to form hydride MH_2 . BeH_2 can be prepared by rxn of BeCl_2 with LiAlH_4



→ **Reducing Nature:** • Weaker reducing agent than corresponding alkali metals.
 • Be has rather less negative reduction potential than other alkaline earth metals but its reducing nature is due to large hydration energy associated with small size of Be^{2+} ion and relatively large value of the atomization enthalpy of the metal.

→ **Solⁿ in liquid NH_3** • dissolves in liq NH_3 to give deep blue black solution forming ammoniated ions.

$$[M + (x+y)\text{NH}_3] \rightarrow [M(\text{NH}_3)_x]^{2+} + 2[e(\text{NH}_3)_y]^-$$
 From these solⁿ, the ammoniates $[M(\text{NH}_3)_6]^{2+}$ can be removed.

USES • Be is used in manufacture of alloys. • Copper-beryllium alloys are used in the prep of high strength springs.
 • Metallic Be is used for making windows X-ray tubes. • Mg forms alloys with Al, Zn, Mn and Sn.
 • Mg-Al alloys being light are used in air-craft construction. • Mg (Powder/Ribbon) is used in flash powders and bulbs, incendiary bombs and single signals. • Suspension of Mg(OH)₂ in H₂O (Milk of Magnesia) used as Antacid.
 • MgCO_3 is used in toothpaste. • Ca is used in extraction of metals from oxides which are difficult to reduce with C.
 • Ca and Ba metal used to remove air from vacuum tubes. • Ra salts are used in radiotherapy (Treatment of Cancer).

COMPOUNDS OF ALKALINE EARTH METAL

→ **Oxides & Hydroxides:** • Burn in O_2 to form MO (monoxide). • Except BeO, they have rock salt structure.
 specially covalent in nature

- Oxides have quite high enthalpies of formation, are quite very stable to heat. • BeO is amphoteric, others are ionic.
- All oxides except BeO are basic, form sparingly soluble hydroxides reacting with H₂O $[MO + H_2O \rightarrow M(OH)_2]$
- Solubility, thermal stability and basic character increase with increase in At. no from Mg(OH)₂ to Ba(OH)₂
- Less basic and less stable than Grp.1 • Be(OH)₂ being amphoteric
$$\begin{aligned} \text{Be(OH)}_2 + 2\text{OH}^- &\rightarrow [\text{Be(OH)}_4]^{2-} \text{ (Beryllate ion)} \\ \text{Be(OH)}_2 + 2\text{HCl} + 2\text{H}_2\text{O} &\rightarrow [\text{Be(OH)}_4]\text{Cl}_2 \end{aligned}$$

→ **Halides:** • Except BeX₂, all halides of Grp.2 are ionic in nature. • BeX₂ is covalent and soluble in organic solvents.
 • BeCl₂ has a chain structure in solid state $[\text{Be} \xrightarrow{\text{Cl}} \text{Be} \xrightarrow{\text{Cl}} \text{Be} \xrightarrow{\text{Cl}} \text{Be} \xrightarrow{\text{Cl}} \text{Be}]$ • In vapour phase, it forms Chloro-bridged dimer which dissociates at order of 1200K. • Tendency to form halide hydrates gradually decreases down $[\text{MgCl}_2 \cdot 8\text{H}_2\text{O}, \text{CaCl}_2 \cdot 6\text{H}_2\text{O}, \text{SrCl}_2 \cdot 6\text{H}_2\text{O}, \text{BaCl}_2 \cdot 2\text{H}_2\text{O}]$
 • Inted of dehydration hydrated halides of Be and Mg on heating suffer hydrolysis.
 • Fluorides are relatively less soluble than chlorides owing high lattice energies.

Salts of Oxacids

- **Carbonates:** • Insoluble in water, can be precipitated by addition of a sodium or ammonium carbonate soluble salts of these metals.
 • The solubility of carbonates decreases with increase in at. no. • All the carbonates decompose on heating to give CO₂ and oxide.
 • BeCO₃ is unstable and kept in CO₂ atmosphere. • Thermal stability increases with increasing cationic size. Thus soluble
- **Sulphate:** • All white solids and stable to heat. • BeSO₄, and MgSO₄ are readily soluble in water.
 • Solubility decrease from CaSO₄ to BaSO₄. • In Be²⁺ and Mg²⁺ ions overcome hydration enthalpies overcome lattice energy enthalpy
- **Nitrates:** • Made by dissolution of the of the carbonates in dil nitric acids. • Mg(NO₃)₂ crystallises with 6 H₂O molecules, Ba(NO₃)₂ crystallises as anhydrous salt. • All decompose on heating $2\text{M(NO}_3)_2 \rightarrow 2\text{MO} + 4\text{NO} + \text{O}_2$

Anomalous Behaviour of Be: [Diagonal relationship with Al]

- Exceptionally small atom and ionic size, Due to high ionisation enthalpy and small size it forms largely covalent compounds
- Due to absence of d orbital, does not exhibit coordination no more than four. Other members can have coordination no of 6.
- Unlike other member, oxides and hydroxide of Be are amphoteric. Easily hydrolysed

Diagonal Relationship b/w Be and Al: • Be²⁺ and Al³⁺ have similar charge / radius ratio

- Al and Be are not readily attacked by acids because of the presence of an oxide film on the surface of the metal.
- Be(OH)₂ dissolved in excess of alkali to give Beryllate ion $[\text{Be(OH)}_4]^{2-}$
- Chlorides of Be and Al have Cl⁻ bridged chloride structure in vapour state. Both chlorides are nonpolar, strong Lewis base. Used as Friedel Craft catalysts.
- Be²⁺ and Al³⁺ have strong tendency to form complexes, BeF_4^{2-} , AlF_6^{3-} . CO₂ is removed quickly.

COMPOUNDS OF Calcium

→ **Quick Lime (CaO):** • Prep'd by heating lime stone (CaCO₃) in a rotary kiln at 1070 - 1270K $[\text{CaCO}_3 \xrightarrow{\text{heat}} \text{CaO} + \text{CO}_2]$

- CaO is a white amorphous solid, mp = 2870K. On exposure to air, it absorbs moisture and carbon dioxide.
- $[\text{CaO} + \text{H}_2\text{O} \rightarrow \text{Ca(OH)}_2 / \text{CaO} + \text{CO}_2 \rightarrow \text{CaCO}_3]$ • Addition of limited water breaks lump of lime, its called slaking of lime
- Quick lime slaked with soda is known as sodalime • Being basic oxide, it combines with acidic oxides at high Temp.
- $[\text{CaO} + \text{SiO}_2 \rightarrow \text{CaSiO}_3 / 6\text{CaO} + \text{P}_4\text{O}_{10} \rightarrow 2\text{Ca}_3(\text{PO}_4)_2]$


→ **Uses:** • For manufacturing cement and is the cheapest form alkali • Manufacturing of sodium carbonate (Na₂CO₃) and (NaOH),
 • In purification of sugar and in manufacture of dye stuffs

→ **Slaked Lime, Ca(OH)₂:** • Prepared by adding CaO to H₂O, white amorphous powder, sparingly soluble in water.


- The aqueous solution is known as lime water and a suspension of Ca(OH)₂ in H₂O is known as milk of lime.
- On passing CO₂, it turns milky due to formation of CaCO₃ $[\text{Ca(OH)}_2 + \text{CO}_2 \rightarrow \text{CaCO}_3 + \text{H}_2\text{O}]$
- On passing excess CO₂, the ppt dissolves to form Ca(HCO₃)₂ $[\text{CaCO}_3 + \text{CO}_2 + \text{H}_2\text{O} \rightarrow \text{Ca(HCO}_3)_2]$ Bleaching Powder
- Milk of lime reacts with Cl₂ to form hypochlorite, a constituent of bleaching powder $[\text{Ca(OH)}_2 + 2\text{Cl}_2 \rightarrow \text{CaCl}_2 + \text{Ca(OCl)}_2 + 2\text{H}_2\text{O}]$
- **Uses:** • preparation of mortar, a building material. • used in white ash due to its disinfectant nature.
- Used in glass making, in tanning industry, prep of bleaching powder and purification of sugar

→ Calcium Carbonate (CaCO_3): CaCO_3 occurs in nature in several forms like lime stone, chalk, marble etc.

• Can be prepared by passing CO_2 through Ca(OH)_2 or by addition of Na_2CO_3 to CaCl_2



• CaCO_3 is a white fluffy powder, almost insoluble in water. When heated at 1200K, it decomposes.



• Reacts with dilute acid to liberate CO_2 $[\text{CaCO}_3 + 2\text{HCl} \rightarrow \text{CaCl}_2 + \text{H}_2\text{O} + \text{CO}_2 / \text{CaCO}_3 + \text{H}_2\text{SO}_4 \rightarrow \text{CaSO}_4 + \text{H}_2\text{O} + \text{CO}_2]$


→ Uses: • Used as building material in form of marble and manufacture of quick lime.

• CaCO_3 and MgCO_3 is used as a flux in the extraction of metals such as iron

• Specially ppt. CaCO_3 is used in making high quality paper.

• Used as antacid, mild abrasive in tooth paste, a constituent of chewing gum and filler in cosmetics.

→ Calcium Sulphate (Plaster of Paris) ($\text{CaSO}_4 \cdot \frac{1}{2}\text{H}_2\text{O}$): • Hemihydrate of CaSO_4 , obtained by heating gypsum, $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ at 393K



• Above 393K, no water of crystallisation is left and anhydrous calcium sulphate, CaSO_4 is formed (dead burnt plaster)

• Has remarkable property of setting with water. On mixing with adequate quantity of water it forms plastic mass that gets into a hard solid in 5 to 15 minutes.

→ Uses: • In building industry as plaster. • Used to immobilise the affected part of organ where there is a bone fracture or sprain, also employed in dentistry, in ornamental works and making casts of statues and busts.

• Cement: Cement is a product obtained by combining a material rich in lime, CaO with other materials such as clay which contains silica, SiO_2 along with the oxides of aluminium.

• The raw materials for the manufacture are limestone and clay, which are mixed together to form 'cement clinker', which is mixed with 2-3% by weight of gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$) to form cement.

• Ingredients of Portland cement (i) dicalcium silicate (Ca_2SiO_4) 26% (ii) tricalcium silicate (Ca_3SiO_5) 51% and (iii) tricalcium aluminate ($\text{Ca}_3\text{Al}_2\text{O}_6$) 11%.

• Setting of Cement: When mixed with water, it settles to a hard mass. This is due to the hydration of the molecules of the constituents and rearrangements.

• The purpose of adding gypsum is only to slow down the process of setting of the cement so that it gets sufficient hardening.

• Biological Importance of Mg and Ca:

• An adult body contains 25g of Mg and 1200g of Ca. • All enzymes that utilise ATP in phosphate transfer require Mg as a cofactor. • Main pigment for the absorption of light in plants is chlorophyll which contains Mg. • About 99% of body Ca is present in bones and teeth, it also plays important roles in neuromuscular function, inter-neuronal transmission, cell membrane integrity and blood coagulation. • Ca concentration in blood plasma is regulated at about 100 mgL^{-1} , it is maintained by two hormones: calcitonin and parathyroid.

P-Block

Group 15 (N, P, As, Sb, Bi) $[ns^2np^3]$

elements: → down the group, there is a shift from non-metallic to metallic through metalloid character. N, P are non metals, As, Sb is metalloids and Bi is a typical metal.

Occurance: → N_2 is 78% of the atmosphere by volume. it occurs in earth's crust as $NaNO_3$, KNO_3 . It is also found in proteins.

→ P appears in members of the apatite family $[Ca_5(PO_4)_3 \cdot CaX_2 (X = F, Cl \text{ or } OH)]$ it is an essential constituent of animal and plant matter (bones, cells).

→ As, Sb and Bi are found mainly as sulphide minerals.

atomic and ionic radii: → increase down the group. considerable increase from N to P and minute in As to Bi (due to d and f orbitals)

ionisation enthalpy: → decreases down the group. due to half filled p orbital, IE is considerably high than group 14. ($\Delta H_1 < \Delta H_2 < \Delta H_3$)

electronegativity: → decreases down the group, not profound in heavier element

physical properties: → N_2 is ~~solid~~ gas, others are diatomic solids. metallic character increases down the group. → b.p increases down the group.

→ m.p. increases upto As then decreases upto Bi.

→ all other than N show allotropy.

CHEMICAL PROPERTIES


ox states and trends in chemical reactivity: → common oxidation states are -3, +3, +5. the tendency to exhibit -3 ox. state decreases down the group. Bi almost doesn't.

→ stability of +5 ox state decreases down the group. only BiF_5 compound is BiF_5 . stability of +3 ox state increases down the group (due to inert pair effect)

→ N also exhibits +1, +2, +4 ox. states reacting with O_2

→ P shows +1 and +4 in some oxoacids

→ all ox states of N from +1 to +4 disproportionate in acid solution.



→ similarly in P, nearly all ox states disproportionate in alkali and acid.

→ N is restricted to a maximum covalency of 4 since only 4 ($1s + 3p$) orbitals are available. others with d orbital can expand their octate.

anomalous properties of nitrogen: → due to small size and high electronegativity, high ionisation enthalpy and non-availability of d orbitals. N has unique ability to form pπ-pπ multiple bond with itself and other elements with small size and high electronegativity (e.g. C, O). thus N_2 exist with high bond dissociation enthalpy.

→ P-P, As-As and Sb-Sb form single bond, Bi forms metallic bond. single N-N is weaker than P-P because of high inter electronic repulsion of the non-bonding electrons. As a result, catenation tendency is weaker in N.

→ due to absence of d orbital in valence shell, it cannot form dπ-pπ ~~per~~ bond as the others. P and As can form dπ-dπ bond also with transition metals when their compounds like $P(C_2H_5)_3$ and $As(C_6H_5)_3$ act as ligands.

reactivity towards H_2 : → all form hydride of form EH_3 . stability of the hydrides decreases from NH_3 to BiH_3 . the reducing nature of the hydrides increases down the group.
→ basicity order; $NH_3 > PH_3 > AsH_3 > SbH_3 > BiH_3$


reactivity towards O_2 : → all form two types of oxides: E_2O_3 and E_2O_5 . oxide with higher ox. state is more acidic. acidic character decreases down the group.
→ E_2O_3 of N, P are purely acidic, As, Sb are amphoteric, Bi is basic.

reactivity towards halogens: → these form EX_3 and EX_5 . N doesn't form NX_5 due to absence of d-orb. → EX_5 are ~~Ex~~ more covalent than EX_3 . except NX_3 all trihalides are stable (NF_3 is stable though) (~~Ex~~ EX_3 except BiF_3 are covalent).


reactivity towards metals: → all form binary compounds exhibiting -3 ox state e.g. Ca_3N_2 , Ca_3P_2 ...



preparation: → produced commercially by liquification and fractional distillation of air
→ treating an aqueous solⁿ of NH_4Cl with sodium nitrite ($NaNO_2$)




→ by thermal decomposition of $(NH_4)_2Cr_2O_7$ [ammonium dichromate]




→ very pure N_2 by thermal decomposition of sodium or barium azide.




properties: → has two stable isotopes: ^{14}N and ^{15}N . very less soluble in water. low freezing and boiling points. rather inert at room temp.
→ reactivity increases rapidly with temperature. at high Temp, it directly combines with some metals to form predominantly ionic nitrides and covalent ones with non-metals.




→ it combines with H_2 at $500^\circ C$ in presence of catalyst (Haber's process)





→ N_2 combines with O_2 at about 2000K to form NO.




preparation: → in nature, due to decay of nitrogenous compounds e.g. urea.




→ in small scale, ammonium salts decompose when treated w/ $NaOH$, $Ca(OH)_2$






→ On large scale, it is manufactured by Haber's process.



properties: → colourless, pungent odour. possesses hydrogen bonds causing high b.p.
→ highly soluble in water, in aq solⁿ is weakly basic due to form OH^-




→ It precipitates the hydroxides of many metals from their salt. e.g.



$$\text{AgCl (s)} + 2\text{NH}_3 \text{ (aq)} \longrightarrow [\text{Ag}(\text{NH}_3)_2] \text{Cl (aq)}$$




EXAMSRoad

→ Zn reacts with dilute nitric acid to give NO_2 and with conc acid to give NO




→ some metals (e.g. Cr, Al) do not dissolve in conc nitric acid because of the formation of a passive film of oxide on the surface.

→ conc HNO_3 also oxidises non-metals and other compounds.



→ the familiar 'Brown Ring Test' for nitrates depends on the ability of Fe^{2+} to reduce nitrates to nitric oxide, which reacts with Fe^{2+} to form a brown coloured complex. the test is usually carried out by adding dilute ferrous sulphate solution to an aqueous solution containing nitrate ion, and then carefully adding concentrated sulphuric acid along the sides of the test tube.




brown



Allotropic forms:

white phosphorus: → translucent white, waxy, poisonous, insoluble in water but soluble in CS_2 and glows in dark.



→ It dissolves in boiling NaOH solution in an inert atmosphere giving PH_3 .



→ it is less stable and therefore, more reactive than others because of angular strain in the P_4 molecule (angles 60°). It readily catches fire in air to give dense white fumes of P_4O_{10} .




red phosphorus:

→ obtained by heating white phosphorus at 573K in inert atmosphere for several days. when it is heated under high pressure, a series of phases of black P is formed.

→ Red P possesses iron grey lustre. odourless, non-poisonous and insoluble in water as well as in ~~carbon~~ CS_2 . Red P is less reactive than white P. doesn't glow in the dark.



→ It is polymeric.




black phosphorus: → it has two forms e.g. α -black and β -black. α -black phosphorus is formed when red P is heated in a sealed tube at 803K. it can be sublimed in air and has opaque monoclinic or rhombohedral crystals. does not oxidise in air,
→ β -black P is prepared by heating white P at 473K under high P . It does not burn in air upto 673K.

Phosphine


preparation: → by the reaction of Calcium phosphide (Ca_3P_2) with H_2O or dilute HCl



→ lab prep: by heating white P with conc $NaOH$ in inert atmosphere of CO_2 .




→ when pure, it is non inflammable but becomes inflammable owing to the presence of P_2H_4 of P_4 vapours. to purify it from impurities, it is absorbed in HI to form phosphonium iodide (PH_4I) which on treating with KOH gives PH_3 .



properties: → colourless gas, rotten fish smell and is highly poisonous. it explodes in contact with traces of oxidising agent like HNO_3 , Cl_2 and Br_2 vapours.

→ slightly soluble in water. solution of PH_3 in water decomposes in presence of light giving red P and H_2 .

→ when absorbed in $CuSO_4$ or $HgCl_2$ solution, the corresponding phosphides formed



→ phosphine is weakly basic and like ammonia, gives phosphonium compounds with acids e.g. $PH_3 + HBr \longrightarrow PH_4Br$


Phosphorus Halides

[PX_3] preparation: → passing dry Cl over heated white P . $P_4 + 6Cl_2 \longrightarrow 4PCl_3$


→ action of thionyl chloride ($SOCl_2$) with white P .



properties: → colourless oily liquid and hydrolyses in the presence of moisture.





→ reacts with organic compounds containing $-OH$ group.



[PX_5] preparation: → reaction of white P with excess of dry chlorine. $P_4 + 10Cl_2 \longrightarrow 4PCl_5$

→ by action of SO_2Cl_2 on P . $P_4 + 10SO_2Cl_2 \longrightarrow 4PCl_5 + 10SO_2$

ClC(Cl)(Cl)Cl
$$2\text{Ag} + \text{PCl}_5 \longrightarrow 2\text{AgCl} + \text{PCl}_3 \quad / \quad \text{Sn} + 2\text{PCl}_5 \longrightarrow \text{SnCl}_4 + 2\text{PCl}_3$$

→ it has trigonal bipyramidal structure. In solid state, it exists as an ionic solid $[PCl_4]^+ [PCl_6]^-$ in which the cation, $[PCl_4]^+$ → tetrahedral, $[PCl_6]^-$ → octa.


Oxoacids of Phosphorus.

$$\begin{array}{c} \text{O} \\ || \\ \text{H} - \text{P} - \text{O} - \text{H} \\ | \\ \text{H} \end{array}$$
OP(=O)(O)O
$$\text{H}_2\text{O} - \text{P}(=\text{O})(\text{OH})_2 - \text{O} - \text{P}(=\text{O})(\text{OH})_2 - \text{OH}$$
$$\text{HO}-\text{P}(=\text{O})(\text{OH})-\text{P}(=\text{O})(\text{OH})-\text{OH}$$
$$\begin{array}{c} \text{OH} \\ | \\ \text{HO} - \text{P} - \text{OH} \\ | \\ \text{OH} \end{array}$$
$$\text{HO}-\overset{\overset{\text{O}}{\parallel}}{\underset{\underset{\text{OH}}{|}}{\text{P}}}-\text{O}-\overset{\overset{\text{O}}{\parallel}}{\underset{\underset{\text{OH}}{|}}{\text{P}}}-\text{OH}$$
O=P(O)(O)OP(=O)(O)OP(=O)(O)OO=P(O)(O)OP(=O)(O)OP(=O)(O)O

cyclotri metaphosphoric acid

poly metaphosphoric
acid.

→ acids with P-H bond have strong reducing properties e.g. H_3PO_2 w/ 2 P-H bonds.



Group 16

elements: O, S, Se, Te, Po electronic configuration: $ns^2 np^4$

atomic and ionic radii: increases down the group. Oxygen is exceptionally small.

ionisation enthalpy: increases down the group. IE of grp 16 is lower compared to grp 15.

electron gain enthalpy: because of compact nature, O atom has less negative electron gain enthalpy than S. from S onwards the value again becomes less negative upto Po.

electronegativity: next to F, O has the highest EN value amongst the elements. EN decreases down the group. metallic character increases from O to Po.

physical properties: \rightarrow O, S are non metals, Se, Te are metalloids, Po is a metal.

\rightarrow Po is radioactive and short lived.

\rightarrow all these elements exhibit allotropy

\rightarrow mp and bp increase with atomic number. large difference b/w the mp and bp of

O and S may be explained on the basis of atomicity [O_2 diatomic, S₈ polyatomic]

CHEMICAL PROPERTIES

oxidation state & chemical reactivity: \rightarrow stability of -2 ox. state decreases down the group. Po doesn't. O only shows -2. exception: OF_2 (+2)

\rightarrow others show +2, +4, +6 ox states. +4 and +6 are more common. S, Se, Te generally show +4 with oxygen and +6 with F. stability of +6 oxidation state decreases down the group and that of +4 increases (inert pair effect).

anomalous behaviour of oxygen: \rightarrow due to small size, high electronegativity. due to this H-bond exists in H_2O but not H_2S .

\rightarrow absence of d orbital limits its covalency to 4 and in practice, it rarely exceeds 2.

reactivity w/ hydrogen: \rightarrow all form hydrides of the type H_2E ($E = O, S, Se, Te, Po$), acidic character increases down

\rightarrow the increase in acidic character down the group can be explained in terms of decrease in bond enthalpy for dissociation of H-E bond down the group.

\rightarrow the thermal stability of hydrides also decreases from H_2O to H_2Po due to this.

\rightarrow all hydrides except water possesses reducing property and this character increases from H_2S to H_2Te .

reactivity w/ oxygen: \rightarrow all form oxides of EO_2 and EO_3 type. O_3 and SO_2 are gases while SeO_2 is solid.

\rightarrow reducing property of dioxide decreases from SO_2 to TeO_2 . SO_2 is reducing while TeO_2 is an oxidising agent.

\rightarrow besides EO_2 , S, Se and Te form EO_3 . both are acidic in nature.


reactivity w/ halogens: \rightarrow they form a large number of halides of type EX_6 , EX_4 and EX_2 .

\rightarrow stability of halides decreases in the order $F^- > Cl^- > Br^- > I^-$. among EX_6 , EX_4 are only stable halides. all EF_6 are gaseous in nature. SF_6 is exceptionally stable for steric reasons.

\rightarrow among EX_4 , SF_4 is a gas, SeF_4 a liquid, TeF_4 a solid.


\rightarrow all except O forms dichlorides and dibromides.

\rightarrow the well known monohalides are dimeric in nature eg. S_2F_2 , S_2Cl_2 , S_2Br_2 , Se_2Cl_2 . these undergo disproportionation as given below.







Dioxygen


preparation: → lab preps: by heating O-containing salts such as chlorates, nitrates and permanganates.



→ lab prep: by thermal decomposition of the oxides of metals low in the electrochemical series and higher oxides of some metals.







→ lab prep: H_2O_2 is readily decomposed into H_2O and O_2 with finely divided metal and MnO_2 as catalyst.



→ on large scale it can be prepared from water or air via electrolysis.

→ industrially, O_2 is obtained by fractional distillation of liquified air after removing CO_2 and water vapour.


properties:




Simple Oxides: → generally, non-metal oxides are acidic but oxides of some metals in high oxidation state have some acidic character (e.g. Mn_2O_7 , CrO_3 , V_2O_5)

→ generally, metal oxides are basic.

→ some show dual behaviour known as amphoteric oxides. e.g.




→ some oxides neither acidic nor basic known as neutral oxides. e.g. CO , NO , N_2O .



→ too reactive to remain in atmosphere at sea level. at about 20 km, ozone layer is formed.

preparation: → passing dry O_2 through a silent electrical discharge, conversion of O_2 to O_3 (10%) occurs. the product is known as ozonised oxygen.





→ as the conversion is endothermic, it is necessary to use a silent electrical discharge in its preparation to prevent its decomposition.

→ for >10%, a battery of ozonisers can be used and pure ozone can be condensed in a vessel surrounded by liquid oxygen.

properties: → pale blue gas, dark blue liquid and violet-black solid. it has a characteristic smell and in small concentration is harmless. however above 100 ppm, breathing becomes uncomfortable resulting in headache and nausea.

→ ozone is thermodynamically unstable wrt O_2 since in decomposition ΔH is -ve and ΔS is positive. $\Delta G < 0$. thus high concentration of O_3 is explosive.

→ due to tendency to liberate atoms of nascent oxygen ($\text{O}_3 \rightarrow \text{O}_2 + \text{O}$), it acts as a powerful oxidising agent. e.g.



→ nitrogen oxides (particularly nitric acid) combine rapidly with O_3 .




→ freons used in aerosol sprays and as refrigerants damages ozone layer.

Sulphur

→ forms numerous allotropes of which yellow rhombic (α -sulphur) and monoclinic (β -sulphur) forms are most important. rhombic sulphur is stable at room temperature.


rhombic sulphur: (α -sulphur) → yellow in colour, formed on evaporation of solution of roll sulphur in CS_2 .



→ insoluble in water but dissolves to some extent in \odot , $R-OH$ and $R-O-R'$.

→ readily soluble in CS_2 .

monoclinic sulphur (β -sulphur)



→ soluble in CS_2 , prepared by melting rhombic sulphur in dish and cooling, till crust is formed. two holes are made in the crust and the remaining liquid is poured out. On removing the crust, colourless needle shaped crystal of β -sulphur. transforms into α -sulphur below 369K.


→ at 369K, both forms are stable, it is known as transition temperature

→ both forms have S_8 molecules with different structure (puckered and crown)

→ in synthesised cyclo- S_6 , it becomes chair form. above 1000K, S_2 is dominant.

Sulphur Dioxide

preparation: → SO_2 is formed with a little (6-8%) SO_3 when S is burnt in air or oxygen.





→ lab prep: treating a sulphite with dilute sulphuric acid





→ industrially, produced as a by product of roasting of sulphide ores.





properties: → colourless, pungent smelling, soluble in water. forms H_2SO_3 with water.



→ reacts readily with $NaOH$ solution forming sodium sulphite which reacts with more SO_2 forming $NaHSO_3$.




→ reacts with Cl_2 in presence of charcoal (catalyst) to give sulphuryl chloride (SO_2Cl_2). it is oxidised to SO_3 by O_2 .




→ moist SO_2 behaves as a reducing agent. it converts $Fe(III)$ ions to $Fe(II)$ and decolourises acidified $KMnO_4$ solution.




Oxoacids of Sulphur




sulphurous acid
(H_2SO_3)



sulphuric acid
(H_2SO_4)



peroxodisulphuric acid
($H_2S_2O_8$)



Pyrosulphuric acid (Oleum)
($H_2S_2O_7$)

Sulphuric Acid


manufacture: → by the Contact Process which involves 3 steps:

- burning of sulphur or sulphide ores in air to generate SO_2 .
- conversion of SO_2 to SO_3 by the reaction with O_2 with V_2O_5 as catalyst.
- absorption of SO_3 in H_2SO_4 to give Oleum ($\text{H}_2\text{S}_2\text{O}_7$)

(ii) is the key step $[\text{2SO}_2 + \text{O}_2 \xrightarrow{\text{V}_2\text{O}_5} \text{2SO}_3, \Delta_r H^\ominus = -196.6 \text{ kJ mol}^{-1}]$. the reaction is exothermic, reversible and forward reaction leads to decrease in volume. therefore, low T and high P are favourable. (generally 2 bar and 720 K is used) dilution of oleum gives H_2SO_4 of the desired concentration (96-98% pure).



properties:

- colourless, dissolves in water with large evolution of heat. [conc acid into water + stir]
- chemical reactions are due to (i) low volatility (ii) strong acid (iii) strong affinity to water (iv) ability to act as oxidising agent.
- due to low volatility, can be used to make more volatile acids from their salts.



- it is strong dehydrating agent (wet gases can be dried up provided they don't react) it removes water from organic compounds. $\text{C}_{12}\text{H}_{22}\text{O}_{11} \xrightarrow{\text{H}_2\text{SO}_4} 12\text{C} + 11\text{H}_2\text{O}$

- hot concentrated sulphuric acid acts as a moderately strong oxidising agent. (intermediate between phosphoric and nitric acid). H_2SO_4 is reduced to SO_2 .



uses: → mainly in making fertilizers, petroleum refining, manufacture of pigments, paints, detergent industry

Group 17 Elements

occurrence: → F, Cl are fairly abundant. At is radioactive. F is normally present as insoluble fluorides (CaF_2 , Na_3AlF_6 etc) and small quantity is present in soil, river water, bones and teeth of animals.

→ sea water contains chlorides, bromides and iodides of Na, K, Mg and Ca (mainly NaCl)

electronic config → $ns^2 np^5$

atomic and ionic radii: → smallest atomic radii in respective ^{periods} ~~group~~. F is extremely small.
→ atomic and ionic radii increases down the group.

ionisation enthalpy: → very high ionisation enthalpy. decreases down the group.

electron gain enthalpy: → maximum electron gain enthalpy in respective ~~group~~ periods because they need only one electron for octate. becomes less negative down the group
→ negative electron gain enthalpy of F is less than Cl due to small size.

electronegativity: → highly electronegative, decreases down the group. F is most electronegative.

physical properties: → all halogens are coloured. (F_2 yellow, Cl_2 greenish yellow, Br_2 red, I_2 violet).

→ F and Cl react with H_2O , Br and I are sparingly soluble in H_2O . but soluble in various organic solvents such as chloroform, CCl_4 , CS_2 and hydrocarbons.

→ enthalpy of dissociation of F_2 is less than that of Cl_2 (X-X). $\text{F}_2 < \text{Cl}_2 > \text{Br}_2 > \text{I}_2$






CHEMICAL PROPERTIES



ox. state and trend in chemical reactivity: → all halogens exhibit -1 ox state. however Cl, Br and I exhibit +1, +3, +5 and +7 also. higher oxidation state of Cl, Br, I are realised mainly when the halogens combine with small and highly electronegative F or O atoms.

→ all halogens are highly reactive. reactivity decreases down the group.

→ due to high acceptance of one electron, are strong oxidising ^{agent} nature. F is the strongest oxidising halogen, oxidises other halide ions in solution or even in solid state. a halogen oxidises halide ions of higher atomic no.



→ relative oxidising power of halogens can further be illustrated by their reaction with water. F oxidises water to O_2 whereas Cl, Br form hydrohalic and hydrohalous acids. I doesn't react spontaneously, I_2 can be oxidised by O_2



anomalous behaviour of fluorine: → due to small size, high electronegativity, low F-F bond dissociation enthalpy and non availability of d orbital.

→ IE, EN and electrode potential are higher than expected.

→ most reactions of F are exothermic (due to small and strong bond formed with other elements).

→ It forms only one oxoacid while others form a number of them.

→ HF is a liquid (due to H-bonding), while other ~~by~~ HX are gases.

reactivity towards H_2 halogens: → all react to give HX but affinity for H_2 decreases from F to I. HX dissolve in H_2O to form hydrohalic acid.

→ acidic strength: $\text{HF} < \text{HCl} < \text{HBr} < \text{HI}$. stability decreases down the group.

reactivity towards O_2 : → forms many oxides but most are unstable, F forms two oxides OF_2 and O_2F_2 only OF_2 is thermally stable. both are strong fluorinating agents. O_2F_2 oxidises Pu to PuF_6 (used in removing Pu from nuclear fuel)

→ Cl, Br, I form oxides of +1 to +7. decrease in stability in order **$\text{I} > \text{Cl} > \text{Br}$**

→ higher oxides tend to be more stable.

→ Cl oxides, Cl_2O , ClO_2 , Cl_2O_6 and Cl_2O_7 are highly reactive ox. agent and tend to explode. Cl_2O is used as bleaching agent for paper pulp, textiles, water treatment

→ Br oxides, Br_2O , BrO_2 , BrO_3 are least stable ~~stable~~ halogen oxides.

→ I oxides, I_2O_4 , I_2O_5 , I_2O_7 are insoluble solids and decompose on heating. I_2O_5 is a very good ox. agent and is used in estimation of CO.


reactivity towards metals: → forms metal halides e.g. $\text{Mg} + \text{Br}_2 \longrightarrow \text{MgBr}_2$

→ ionic character $\text{MF} > \text{MCl} > \text{MBr} > \text{MI}$. ~~same~~ metals with higher ox. state is more stable and more covalent.


reactivity towards halogens: → they combine to form compounds of types XX' , XX'_3 , XX'_5 and XX'_7 , where X is a larger size and X' is smaller size halogens.

Chlorine


Preparation: → heating MnO_2 with concentrated HCl . $MnO_2 + 4HCl \longrightarrow MnCl_2 + Cl_2 + 2H_2O$
however $NaCl$ and conc H_2SO_4 is used in stead of HCl .



→ by the action of HCl on $KMnO_4$.





manufacture: → Deacon's process: oxidation of HCl gas by atmospheric O_2 with $CuCl_2$ cat. at ~~723~~ 723K.



→ Electrolytic process: Cl_2 is obtained by electrolysis of brine (conc. $NaCl$ solution). Cl_2 is liberated at anode.

properties: → greenish yellow gas with pungent and suffocating odour. It is about 2-5 times heavier than air. It is 2-5 times heavier than air. can be easily liquified.

→ reacts with metals and non-metals to form chlorides.





→ It has great affinity for hydrogen, reacts with H -compounds forming HCl




→ with excess ammonia, Cl_2 gives N_2 and NH_4Cl but with excess Cl_2 , NCl_3 (explosive) is formed.





→ with cold and dilutes, Cl_2 produces a mixture of chloride and hypochlorite but not with hot and concentrated alkalis it gives chloride + chlorate.



→ with dry slaked lime it gives bleaching powder $[Ca(OCl)_2 \cdot CaCl_2 \cdot Ca(OH)_2 \cdot 2H_2O]$







→ Cl_2 reacts with hydrocarbons and gives substitution products with saturated hydrocarbons and addition products with unsaturated hydrocarbons.





→ Chlorine water on standing loses its yellow colour due to the formation of HCl and $HOCl$. Hypochlorous acid ($HOCl$) so formed, gives nascent oxygen which is responsible for oxidising and bleaching properties of Cl_2 .

→ it oxidises ferrous to ferric and sulphite to sulphate, sulphur to SO_3 , iodine to iodate. In presence of H_2O , they form H_2SO_4 and HIO_3 .





→ It is a powerful bleaching agent; bleaching action is due to oxidation.




Hydrogen Chloride

preparation: lab: by heating NaCl with concentrated sulphuric acid.




properties: → bleaches veg and organic matter in presence of moisture. bleaching effect of Cl_2 is permanent.

→ colourless and pungent smelling gas, easily liquified and extremely soluble in water and ionises.





→ its aqueous solution is called hydrochloric acid.

→ when 3 parts of conc HCl and one part of concentrated HNO_3 are mixed, Aqua Regia is formed which is used for dissolving noble metals e.g. gold, platinum



→ decomposes salts of weaker acid e.g. carbonates, hydrogencarbonates, sulphites etc.



Oxoacids of Halogens

→ due to high EN and small size, F forms only one oxoacid. HOF known as fluoric (I) acid or hypofluorous acid. Others form several oxoacids which cannot be isolated in pure form. they are stable in their salts and in aqueous solution.

Halic (I) acid (hypohalous)

HOF
(hypofluorous)

HOCl
(hypochlorous)

HOBr
(hypobromous)

HOI
(hypoiodous)

Halic (III) acid (Halous)

HOClO
(chlorous)

HOBrO₂
(bromic)

HOIO₂
(iodic)

Halic (V) acid (Halic)

HOClO₂
(chloric acid)

HOBrO₃
(perbromic)





HOIO₃
(periodic)

Halic (VII) acid (Perhalic)

HOClO₃
(perchloric)







HOBrO₃
(perbromic)

HOIO₃
(periodic)




INTERHALOGEN COMPOUNDS

preparation: → by direct combination or by the action of halogen on lower interhalogen compounds.



properties: → covalent molecules and diamagnetic in nature. generally more reactive than halogens (except F) as $\text{X}-\text{X}'$ bond is weaker than $\text{X}-\text{X}$ (except $\text{F}-\text{F}$).


→ all undergo hydrolysis giving halide ion derived from the smaller halogen and a hypohalite (when XX'), halite (XX'_3), halate (when XX'_5) and perchalate (when XX'_7)



GROUP 18 ELEMENTS

occurrence: → all except Rn occur in the atmosphere. He and sometimes Ne are found in minerals of radioactive origin e.g. pitchblende, monazite, cleveite.

→ commercial source of helium is natural gas. Rn is obtained as a decay product of ^{226}Ra



electron configuration: → ns^2np^6 except helium which is $1s^2$.

atomic radii: → increases down the group

ionisation enthalpy: → exhibit very high ionisation enthalpy. decreases down the group.




electron gain enthalpy: → no tendency to accept electron therefore, have large positive values of EGA.

physical properties: → all monoatomic, colourless, odourless and tasteless. They are sparingly soluble in H_2O . very low m.p and b.p. He has the lowest b.p. (4.2K) of any known substance.
→ it has unusual property of diffusing through most commonly used lab materials such as rubber or plastics.

Chemical properties: → there are several Xe compounds, compounds of Kr are fewer. only KrF_2 has been studied in detail.

→ Compounds of Rn has not been isolated just identified (e.g. RnF_2).


Xenon-F compounds: → forms XeF_2 , XeF_4 and XeF_6 but combination under appropriate experimental conditions.






→ XeF_6 can also be prepared by interaction of XeF_4 and O_2F_2 at 143K





→ XeF_2 , XeF_4 and XeF_6 are colourless crystalline solids and sublime readily at 25°C. they are powerful fluorinating agents. readily hydrolysed even by traces of H_2O .



→ Xe fluorides react with F ion acceptors to form cationic species and fluoride ion donors to form fluoranions.



Xenon-O compounds: → Hydrolysis of XeF_4 and XeF_6 with H_2O gives XeO_3 .



→ partial hydrolysis of XeF_6 gives oxyfluorides, XeOF_4 and XeO_2F_2

→ XeO_3 is a colourless explosive solid and has a pyramidal molecular structure.

→ XeOF_4 is a colourless volatile liquid and has a square pyramidal molecular structure.